

**HUBUNGAN POLA MAKAN DENGAN STATUS GIZI PENDERITA
TB PARU DI WILAYAH PUSKESMAS SUKA MAKMUR DAN
PUSKESMAS SEBLAT BENGKULU UTARA
TAHUN 2018**

SKRIPSI



OLEH

MIDIAWATI

**PROGRAM STUDI ILMU KEPERAWATAN (S-1) FAKULTAS
ILMU KESEHATAN (FIKES) DEHASEN BENGKULU
UNIVERSITAS DEHASEN BENGKULU
TAHUN 2018**

ABSTRACT

Tuberculosis is an infectious disease which cause by Mycobacterium Tuberculosis bacteria infection and still becoming public health problem where there are a million people died because of TB. One effect which occurred on Pulmonary TB patient is abnormal nutritional status. The purpose of this research is to find out Correlation between eating pattern with nutrition status of Pulmonary Tuberculosis patient at SukaMakmur Public Health and Seblat Public Health North Bengkulu in 2018.

The method used in this research is analytic method, used cross sectional design. Population in this research is 43 Pulmonary TB patients at SukaMakmur public health and seblat public health in 2017 an 43 samples data are taken with total sampling. Data analysis which used in this research is using univariate and bivariate using chi square analysis.

Univariate analysis result shows that almost half of respondents (51,2%) have bad eating pattern and half respondents)55,8%) have normal nutritional status, while according to chi-square analysis shows that $p=0,002$ smaller than alpha 0,05. It means there is a correlation between eating pattern with nutritional status of pulmonary TB.

It is expected that to all public health institution can improe observation and counseling to improve the knowledge about Pulmonary TB especially the important of good eating pattern thru improvement of nutrition counseling.